

**EVALUATION OF PERFORMANCE AND CARCASS
CHARACTERISTICS OF BROILER CHICKEN FED GRADED LEVELS OF
RAW COCOA (*Theobroma cacao* lin) BEAN SHELL BASED DIET.**

M.D OLUMIDE, R.A HAMZAT, O.A AJAYI*

Department of Animal Science

**School of Agriculture and Industrial Technology Babcock
University ilishan Remo,**

Ogun state.

**Department of Animal Science, Federal University Dutsin-Ma,
Dutsin-Ma**

kaduna state.

INTRODUCTION

- Poultry production in the tropics is facing the challenges of poor nutrition. (Babayemi *et al*, 2000).
- Livestock and poultry generally compete with humans for maize grain. The competition that exist between man and his livestock has necessitated the search for alternative ingredients.
- Large quantities of cocoa bean shell are produced and wasted annually by farmers and associated processing industries in Nigeria.

- Cocoa bean shell (CBS) is a waste from cocoa processing industries in Nigeria and it constitutes a serious disposal problem (Hamzat,*et.al.*,2006; Olumide, *et al.*, 2008).
- There is paucity of information on the use of graded levels of raw cocoa bean shell in the diets of broiler birds.
- Therefore, the objective of this study is focused on evaluation of performance and carcass characteristics of broilers fed graded levels of cocoa bean shell based diets.

MATERIALS AND METHOD

- A total of one hundred and fifty day old broilers were used for this research. This experiment was carried out at University of Ibadan, Teaching and Research Farm, livestock section for a period of eight weeks.
- The birds were randomly allotted to five dietary treatments with ten birds per replicate in a completely randomized design. The treatments were; A (0 % CBS- control), B (5 % RCBS), C (10 % RCBS), D (15 % RCBS) and E (20 % RCBS). The birds were fed *ad libitum*. Weekly records of body weight and daily feed intake were taken. Feed conversion ratio was calculated as a ratio of feed consumption and bodyweight gain.

- At the end of 8 weeks, five birds were randomly selected from each of the replicates for carcass analysis. The selected birds were starved overnight and their live weights recorded. The birds were slaughtered by severing the jugular vein and were fully bled before scalding in hot water. The birds were de-feathered and their plucked weights were taken.
- Eviscerated dressed, thigh, drumsticks and organ (Liver, heart and gizzard) weight were expressed as percentage of live weight. All data collected were subjected to analysis of variance (ANOVA) of SAS, (1999) and significant means were separated using Duncan multiple range test of the same software.

TABLE 1: Gross Composition of Broiler Starter Containing Graded Level of Raw Cocoa Bean Shell

Ingredients (kg)	A (0%)	B (5%)	C (10%)	D (15%)	E (20%)
Maize	57.00	54.15	51.30	48.45	45.60
Cocoa Bean Shell (CBS)	-	2.85	5.70	8.55	11.40
Soybean meal	33.51	33.51	33.51	33.51	33.51
Fishmeal 72%	1.00	1.00	1.00	1.00	1.00
Full fat Soya	4.10	4.10	4.10	4.10	4.10
Oyster shell	0.50	0.50	0.50	0.50	0.50
Bone meal	3.30	3.30	3.30	3.30	3.30
Methionine	0.04	0.04	0.04	0.04	0.04
Salt	0.30	0.30	0.30	0.30	0.30
Premix	0.025	0.025	0.025	0.025	0.025
Total	100.00	100.00	100.00	100.00	100.00
CALCULATED NUTRIENTS					
Crude Protein (%)	23.21	23.41	23.62	23.83	24.04
Metabolisable Energy (Kcal/kg)	2951.62	2925.57	2899.52	2873.47	2847.40
Crude Fibre (%)	4.01	4.34	4.66	4.99	5.35

Table 2: Gross Composition of Finisher Diet Containing Graded Level of Raw Cocoa Bean shell

Ingredients (kg)	A (0%)	B (5%)	C (10%)	D (15%)	E (20%)
Maize	57.00	54.15	51.30	48.45	45.60
Cocoa Bean Shell (CBS)	-	2.85	5.70	8.55	11.40
Soybean meal	25.80	25.80	25.80	25.80	25.80
Palm kernel oil	2.47	2.47	2.47	2.47	2.47
Full fat Soya	10.00	10.00	10.00	10.00	10.00
Oyster shell	0.50	0.50	0.50	0.50	0.50
Bone meal	3.30	3.30	3.30	3.30	3.30
Methionine	0.08	0.08	0.08	0.08	0.08
Salt	0.30	0.30	0.30	0.30	0.30
Premix	0.025	0.025	0.025	0.025	0.025
Total	100.00	100.00	100.00	100.00	100.00
CALCULATED NUTRIENTS					
Crude Protein (%)	21.48	21.69	21.90	22.11	22.32
Metabolisable Energy (Kcal/kg)	3122.33	3096.28	3070.23	3044.18	3080.13
Crude Fibre (%)	2.81	4.14	4.47	4.80	5.13

Table 3: Proximate Composition of Raw Cocoa Bean Shell

Dry Matter (%)	89.00
Moisture (%)	11.00
Crude Protein (%)	14.58
Crude Fiber (%)	15.58
Ash (%)	9.70
Ether Extract (%)	2.79
Nitrogen Free Extract (%)	57.10
Metabolisable Energy (k cal/kg)	2308.10
Theobromine (%)	0.85

performance characteristics of broiler chicken fed graded level of raw cocoa bean shell

.

- There were significant ($P < 0.05$) difference in feed intake between birds on the control and RCBS supplemented diets. The weight gain and feed conversion ratio of bird fed the RCBS based diets were significantly ($P < 0.05$) different from their contemporary on the control diets
- The mortality rate was highest with the bird fed with diet containing 20 % RCBS. This showed that the tolerable level of raw cocoa bean shell was exceeded at 20%.

Table 4: Performance Characteristics of Broiler Chicken Fed Cocoa Bean Shell

Parameters	A (0%)	B (5%)	C (10%)	D (15%)	E (20%)	SEM
AVERAGE FEED INTAKE {g}	4556.50 ^a	4529.00 ^b	4200.50 ^c	4177.50 ^d	4097.00 ^e	2.33
INITIAL WEIGHT {g}	37.00	37.00	37.00	36.00	36.00	-
FINAL WEIGHT {g}	1952.00 ^a	1812.00 ^b	1739.00 ^c	1630.00 ^d	1550.00 ^e	7.34
AVERAGE WEIGHT GAIN {g}	1915.00 ^b	2076.56 ^a	1901.00 ^b	1806.00 ^c	1665.00 ^d	1.84
FCR	2.37 ^c	2.55 ^b	2.58 ^b	2.40 ^b	2.81 ^a	0.21
MORTALITY %	0.00	3.33	3.33	6.66	10.00	0.01

Table 5: Carcass Characteristics of Broilers Fed Graded Levels of Raw Cocoa Bean Shell Based Diets

PARAMETERS	0%	5%	10%	15%	20%	SEM
	Control	RCBS	RCBS	RCBS	RCBS	
Live weight(g)	1952.00 ^a	1812.33 ^b	1739.67 ^c	1630.67 ^d	1550.67 ^e	9.38
Plucked weight(g)	1752.00 ^a	1708.67 ^a	1489.67 ^b	1380.00 ^b	1360.67 ^c	9.15
Eviscerated weight (g)	1518.67 ^a	1493.33 ^a	1348.67 ^b	1280.33 ^b	1130.00 ^c	8.09
Visceral organ	233.33 ^a	200.00 ^b	150.00 ^d	146.67 ^e	176.67 ^c	8.56
Dressed weight(g) %	1438.03 ^a	1286.88 ^b	1215.50 ^b	1116.67 ^c	1116.67 ^c	12.9
Dressed %	73.67 ^a	71.02 ^a	69.90 ^a	67.00 ^b	69.07 ^a	1.23
Breast %	23.00 ^a	23.00 ^a	22.00 ^a	21.00 ^b	19.80 ^b	2.34
Thigh %	15.76	15.55	15.69	15.36	14.15	2.12
Drumstick %	14.99	14.50	14.50	14.00	14.00	1.52
Organ weight						
Gizzard %	3.24	3.08	3.36	3.41	3.59	0.88
Liver %	2.76 ^d	3.33 ^c	3.88 ^c	4.34 ^b	5.31 ^a	0.40
Heart %	0.68	0.66	0.66	0.62	0.61	0.12

a, b, c, d, e = Means along the same rows with different superscripts are significant (P < 0.05)

RCBS = Raw Cocoa Bean shell; SEM=Standard Error of Means.

Carcass Characteristics of Broiler Fed Raw Cocoa Bean Shell

- From the carcass characteristics of broilers fed raw cocoa bean shell based diets.
- Significant difference ($P < 0.05$) were obtained in the values of plucked weight, eviscerated weight, visceral organ, dressed % while the values obtained for thigh, drumstick, gizzard and heart were not significantly different.

CONCLUSION

- It can therefore be concluded that broiler can effectively utilize RCBS up to 10% inclusion in the diet, without compromising the performance and the health status of the birds.

THANK YOU FOR YOUR AUDIENCE

- **REFERENCES**

- Babayemi, O J, D.Demeyer and V. Fievez (2000). Nutritive Value and Qualitative Assessment of Secondary Compound in Seeds of Eight Tropical Browse, Shrub and Pulse Legumes. Comm. App. Biol. Scientist University, 69/I: 103-110.
- Hamzat, R. A., Babatunde, B. B., Adejinmi, O. O., Olubamiwa, O. (2006). Utilization of cocoa bean shell as a feed ingredient for broiler and cockerel chickens. Paper presented at the 15th International Cocoa Research Conference (ICRC) held at the International Conferences Center, Ramada Plaza, Costa Rica, between 8th and 14th October, 2006.
- Olumide, M .D, Hamzat, R.A, And Akinsoyinu O. A (2008) Carcass Evaluation Of **Finishing** Broilers Fed Fermented Cocoa Bean Shells Based Diets. International Poultry Scientific Forum, Held At World Congress Center, Atlanta, Georgia .On Jan 21-22.
- SAS Institute (1999). SAS START Users Guide ® Statistical Version 9 Edition SAS Institute Inc., Cary, North Carolina, USA.